

CD31 Recombinant antibody

Cat: B03026R

Company: Haokebio

Uniprot ID: Q08481

Applications: IHC:1:250-1:500

Organism: Rabbit

IHC-Polymer:1:1000-1:2000

Species reactivity: Human Mouse

IHC-TSA:1:1200-1:2500

Background:

Platelet endothelial cell adhesion molecule-1 (PECAM-1, CD31) is a member of the immunoglobulin gene superfamily of cell adhesion molecules. CD31 is a transmembrane glycoprotein that is highly expressed on the surface of the endothelium, making up a large portion of its intracellular junctions. PECAM-1 is also present on the surface of hematopoietic cells and immune cells including platelets, monocytes, neutrophils, natural killer cells, megakaryocytes and some types of T-cell. As well as its role in cell-cell adhesion, PECAM-1 functions as a signaling receptor, and is involved in important physiological events such as nitric oxide production, regulation of T-cell immunity and tolerance, leukocyte transendothelial migration and inflammation and angiogenesis.

Protein full name:

platelet/endothelial cell adhesion molecule 1

Synonyms:

Pecam, PECAM 1, Pecam1, PECAM-1

Immunogen:

Recombinant protein

Isotype:

IgG

Subcellular location:

Membrane

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.


Storage:

Store at -20 °C for one year.


Experimental procedure:

Antigen retrieval: Citrate buffer (pH 9.0), Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Images:



Sample: Mouse tumor, 4% PFA 12-24h



Sample: Mouse tumor, 4% PFA 12-24h

Source of Reagents:

发表[中文论文]请标注:CD31(B03026R)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:CD31(B03026R) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.

